

CLINICAL OUTCOMES FOLLOWING PROTON BEAM THERAPY FOR LOCALLY ADVANCED NON-SMALL CELL LUNG CANCER: ANALYSIS OF A MULTI-INSTITUTIONAL PROSPECTIVE REGISTRY

Aurélien Jongen*, Florian Charlier, M.D.*, Luigi Moretti, M.D., Ph.D.*, Dirk Van Gestel, M.D., Ph.D.*, Kelsey Baker†, Mary Redman, Ph.D.†, Carlos Vargas, M.D.‡, Gary Larson, M.D.§, William Hartsell, M.D.¶, Henry Tsai, M.D.¶, George Laramore, M.D.#, Lane Rosen, M.D.**, Mark Mishra, M.D.††, and Ramesh Rengan, M.D., Ph.D.#

*Department of Radiation Oncology, Institut Jules Bordet, Université Libre de Bruxelles, Brussels, Belgium; †Clinical Statistics, Clinical Research Division, Fred Hutchinson Cancer Center, Seattle, Washington; ‡Mayo Clinic Arizona Proton Therapy Program, Rochester, Minnesota; §Oklahoma Proton Center, Oklahoma City, Oklahoma; ¶Northwestern Medicine Chicago Proton Center, Warrenville, Illinois; ¶New Jersey Proton Therapy Center, Somerset, New Jersey; #University of Washington and Seattle Cancer Care Alliance Proton Therapy Center, Seattle, Washington; **Willis-Knighton Medical Center, Shreveport, Louisiana; ††University of Maryland, Baltimore, Maryland;

Background: Level 1 evidence is still lacking for proton beam therapy (PBT). By identifying target populations that would benefit most from PBT, prospective registries could overcome the challenges in clinical trials enrollment. Herein, we report clinical outcomes of patients treated with PBT for locally advanced non-small cell lung cancer (LA-NSCLC).

Method & Materials: Data were obtained from the prospective database of the Proton Collaborative Group (PCG), collecting treatment-related information from twelve PBT centers across the USA. Inclusion criteria of our study were stage III LA-NSCLC, use of PBT, and availability of follow-up data. Survival time was calculated from start of treatment until death or last follow-up. Kaplan-Meier curves were generated for groups of interest and compared with log-rank tests. Cox regression modeling was used to evaluate the relationship between covariates of interest and overall survival (OS).

Results: A total of 195 patients were included in the analysis. PBT alone was given to 93% of patients with a median equivalent dose in 2 Gy fractions (EQD2) of 63.8 Gy(RBE). Pencil beam scanning (PBS) was used in 20% of treatments. Treatment-related grade 3 adverse events (AEs) were rare: one pneumonitis, two dermatitis and three esophagitis. No grade 4 events were reported. Two grade 5 events occurred, both cardiological, probably unrelated to PBT. The median follow-up time for living patients was 13.6 months and the median OS was 19.0 months. On multivariate analysis, good performance status, PBS use, and increased EQD2 showed a statistically significant association with higher OS.

Conclusion: PBT appears to yield low rates of AEs with encouraging OS for the treatment of LA-NSCLC. Use of PBS and increased treatment dose can potentially increase OS. Prospective databases such as the PCG registry could play a key role in the future but need thorough updates to reflect the clinical reality.